

Environmental Protection Agency

§ 180.486

Compliance with the tolerance levels specified below is to be determined by measuring only flutolanil and its metabolites converted to 2-(trifluoromethyl) benzoic acid and calculated as flutolanil, in or on the following commodities.

Commodity	Parts per million
Wheat, bran	0.20
Wheat, forage	2.5
Wheat, grain	0.05
Wheat, hay	1.2
Wheat, straw	0.20

[60 FR 42458, Aug. 16, 1995, as amended at 61 FR 33044, June 26, 1996; 63 FR 42256, 42257, Aug. 7, 1998; 66 FR 10825, Feb. 20, 2001; 71 FR 74818, Dec. 13, 2006; 72 FR 35665, June 29, 2007; 73 FR 33017, June 11, 2008; 75 FR 17570, Apr. 7, 2010; 75 FR 80350, Dec. 22, 2010]

§ 180.485 Cyproconazole; tolerances for residues.

(a) *General.* (1) Tolerances are established for residues of the free and conjugated forms of the fungicide cyproconazole, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the proposed tolerance levels specified below is to be determined by measuring only cyproconazole (α -(4-chlorophenyl)- α -(1-cyclopropylethyl)-1*H*-1,2,4-triazole-1-ethanol) in or on the following commodities:

Commodity	Parts per million
Aspirated grain fractions	2.5
Cattle, fat	0.01
Cattle, meat byproducts (except liver)	0.01
Coffee bean, green (Imported) ¹	0.1
Corn, field, forage	0.60
Corn, field, grain	0.01
Corn, field, stover	1.2
Goat, fat	0.01
Goat, meat byproducts (except liver)	0.01
Horse, fat	0.01
Horse, meat byproducts (except liver)	0.01
Peanut	0.01
Peanut, hay	6.0
Sheep, fat	0.01
Sheep, meat byproducts (except liver)	0.01
Soybean, forage	1.0
Soybean, hay	3.0
Soybean, oil	0.10
Soybean, seed	0.05
Wheat, forage	0.80
Wheat, grain	0.05
Wheat, grain, milled byproducts	0.10
Wheat, hay	1.3
Wheat, straw	0.90

¹There are no U.S. registrations as of February 15, 2008 for use on coffee bean.

(2) A tolerance is established for the combined residues of the free and conjugated forms of the fungicide cyproconazole, including its metabolites and degradates, in or on the commodity in the table below. Compliance with the tolerance level specified below is to be determined by measuring only the sum of cyproconazole (α -(4-chlorophenyl)- α -(1-cyclopropylethyl)-1*H*-1,2,4-triazole-1-ethanol) and its metabolite δ -(4-chlorophenyl)- β , δ -dihydroxy- γ -methyl-1*H*-1,2,4-triazole-1-hexenoic acid, calculated as the stoichiometric equivalent of cyproconazole, in or on the following commodity:

Commodity	Parts per million
Milk	0.02

(3) Tolerances are established for the combined residues of the free and conjugated forms of the fungicide cyproconazole, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the tolerance level specified below is to be determined by measuring only the sum of cyproconazole (α -(4-chlorophenyl)- α -(1-cyclopropylethyl)-1*H*-1,2,4-triazole-1-ethanol) and its metabolite 2-(4-chlorophenyl)-3-cyclopropyl-1-[1,2,4]triazol-1-yl-butane-2,3-diol, calculated as the stoichiometric equivalent of cyproconazole, in or on the following commodities:

Commodity	Parts per million
Cattle, liver	0.50
Goat, liver	0.50
Hog, liver	0.01
Horse, liver	0.50
Sheep, liver	0.50

(b) *Section 18 emergency exemptions.* [Reserved]

(c) *Tolerances with regional registrations.* [Reserved]

(d) *Indirect or inadvertent residues.* [Reserved]

[63 FR 53835, Oct. 7, 1998, as amended at 71 FR 71058, Dec. 8, 2006; 73 FR 27760, May 14, 2008; 78 FR 37473, June 21, 2013]

§ 180.486 Chlorethoxyfos; tolerances for residues.

(a) *General.* Tolerances are established for residues of the insecticide